



VYVGART® Hytrulo

(efgartigimod alfa and
hyaluronidase-qvfc)

Subcutaneous Injection
180 mg/mL and 2000 U/mL vial

VYVGART®
(efgartigimod alfa-fcab)

Injection for Intravenous Use
400 mg/20 mL vial

GENERALIZED MYASTHENIA GRAVIS

*doesn't get to steal
this moment*

Getting started with VYVGART

What is VYVGART® (efgartigimod alfa-fcab) for intravenous (IV) infusion and what is VYVGART® HYTRULO (efgartigimod alfa and hyaluronidase-qvfc) for subcutaneous injection?

VYVGART and VYVGART HYTRULO are both prescription medicines, each used to treat a condition called generalized myasthenia gravis, which causes muscles to tire and weaken easily throughout the body, in adults who are positive for antibodies directed toward a protein called acetylcholine receptor (anti-AChR antibody positive).

SELECT IMPORTANT SAFETY INFORMATION

VYVGART and VYVGART HYTRULO may increase the risk of infection. The most common infections for efgartigimod alfa-fcab-treated patients were urinary tract and respiratory tract infections.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

Generalized myasthenia gravis (gMG) is a rare, autoimmune, neuromuscular condition that causes muscle weakness and fatigue

MUSCLE WEAKNESS FROM gMG CAN CAUSE:

- » Eyelid drooping
- » Blurred or double vision
- » Difficulty speaking
- » Difficulty chewing/swallowing
- » Choking
- » Difficulty supporting neck
- » Shortness of breath/difficulty breathing
- » Weakness in the arms and legs
- » Difficulty walking/standing
- » Fatigue from repeated muscle use

gMG symptoms can make daily life more challenging and may limit one's ability to do everyday activities

UNDERSTANDING HARMFUL AChR ANTIBODIES IS AN IMPORTANT PART OF UNDERSTANDING gMG



Antibodies, also known as immunoglobulins (Ig), are proteins produced by the immune system to help **protect the body from infection and disease.**



The immune system makes **5 different types of antibodies (IgG, IgA, IgE, IgD, IgM).** IgG is the most common.



Sometimes the immune system mistakenly makes harmful IgG antibodies that target AChR. Patients with these harmful AChR antibodies have **anti-AChR antibody positive gMG.**

HARMFUL AChR ANTIBODIES CAUSE gMG SYMPTOMS



While IgG antibodies continue to protect the body, **harmful AChR antibodies disrupt communication between muscles and nerves.**



This **prevents muscles from getting some messages sent by nerves;** this is what causes gMG symptoms.

WITH ANTI-AChR ANTIBODY POSITIVE gMG



➔ **Learn more about harmful AChR antibodies and gMG at [ABOUTgMG.com](https://www.ABOUTgMG.com)**

ACh=acetylcholine;
AChR=acetylcholine receptor;
gMG=generalized myasthenia gravis;
IgG=immunoglobulin G
Visit [VYVGART.com/glossary](https://www.VYVGART.com/glossary) for a glossary of terms.



**GENERALIZED
MYASTHENIA
GRAVIS**
*doesn't get to
interrupt this moment*


Have questions?
Call 1-833-VYVGART (1-833-898-4278)
to speak with an educator

A FIRST-OF-ITS-KIND TREATMENT

VYVGART® Hytrulo
(efgartigimod alfa and hyaluronidase-qvfc)
Subcutaneous Injection
180 mg/mL and 2000 U/mL vial

VYVGART®
(efgartigimod alfa-fcab)
Injection for Intravenous Use
400 mg/20 mL vial

VYVGART is the first FDA-approved treatment that uses a fragment of an IgG antibody to treat adults with anti-AChR antibody positive gMG



VYVGART is a fragment of an IgG antibody.

Choose the VYVGART option that is best for you

VYVGART® Hytrulo

(efgartigimod alfa and hyaluronidase-qvfc)
Subcutaneous Injection
180 mg/mL and 2000 U/mL vial

For **subcutaneous injection**—given by delivering medicine into the layer beneath the skin.

Pages 14 and 15: effectiveness and safety information

VYVGART®

(efgartigimod alfa-fcab)
Injection for Intravenous Use
400 mg/20 mL vial

For **IV infusion**—given by inserting a needle into a vein so the medicine enters the bloodstream.

Pages 10 and 11: effectiveness and safety information

IV=intravenous

Visit [VYVGART.com/glossary](https://www.vyvgart.com/glossary) for a glossary of terms.

SELECT IMPORTANT SAFETY INFORMATION


Before taking VYVGART or VYVGART HYTRULO, tell your doctor if you: take any medicines, including prescription and non-prescription medicines, supplements, or herbal medicines; have received or are scheduled to receive a vaccine (immunization); or have any allergies or medical conditions, including if you are pregnant or planning to become pregnant, or are breastfeeding.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

HOW VYVGART WORKS

Harmful AChR antibodies cause gMG symptoms

Receptors called “FcRn” extend the life of IgG antibodies, including the harmful AChR antibodies that cause gMG symptoms. This allows harmful AChR antibodies to continue causing symptoms. **IgG antibodies, including harmful AChR antibodies, that can’t attach to FcRn are removed by the body.**



SELECT IMPORTANT SAFETY INFORMATION

VYVGART and VYVGART HYTRULO may cause serious side effects, including infection. VYVGART and VYVGART HYTRULO may increase the risk of infection. The most common infections for efgartigimod alfa-fcab-treated patients were urinary tract and respiratory tract infections. Signs or symptoms of an infection may include fever, chills, frequent and/or painful urination, cough, pain and blockage of nasal passages/sinus, wheezing, shortness of breath, fatigue, sore throat, excess phlegm, nasal discharge, back pain, and/or chest pain.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

VYVGART causes IgG antibodies to be reduced, including the harmful AChR antibodies that cause gMG symptoms

VYVGART specifically works to reduce IgG antibodies, including gMG-causing, harmful AChR antibodies. VYVGART targets FcRn receptors, preventing many IgG antibodies, including the harmful AChR antibodies, from attaching. **Harmful AChR antibodies that don't attach to FcRns are removed from the body and can no longer cause gMG symptoms.**




AChR=acetylcholine receptor; FcRn=neonatal Fc receptor; gMG=generalized myasthenia gravis; IgG=immunoglobulin G
Visit [VYVGART.com/glossary](https://www.vyvgart.com/glossary) for a glossary of terms.

SELECT IMPORTANT SAFETY INFORMATION

What are the most common side effects of VYVGART and VYVGART HYTRULO? The most common side effects in efgartigimod-alfa-fcab-treated patients were respiratory tract infection, headache, and urinary tract infection. Additional common side effects with VYVGART HYTRULO are injection site reactions, including rash, redness of the skin, itching sensation, bruising, pain, and hives. These are not all the possible side effects of VYVGART and VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to the US Food and Drug Administration at 1-800-FDA-1088.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).



**GENERALIZED
MYASTHENIA
GRAVIS**

*doesn't get to stop
this cheering section*

VYVGART FOR IV INFUSION



**VYVGART is a 1-hour IV infusion
given at an infusion center,
neurologist's office, or at home***

*In some cases, VYVGART may also be given at home by a trained nurse.

IV=intravenous

Visit [VYVGART.com/glossary](https://www.vyvgart.com/glossary) for a glossary of terms.

SELECT IMPORTANT SAFETY INFORMATION

Do not use VYVGART if you have a serious allergy to efgartigimod alfa or any of the other ingredients in VYVGART. Do not use VYVGART HYTRULO if you have a serious allergy to efgartigimod alfa, hyaluronidase, or any of the other ingredients in VYVGART HYTRULO. VYVGART and VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

EFFECTIVENESS

When added to their current gMG treatment, VYVGART helped patients in the study with anti-AChR antibody positive gMG achieve:

Improved daily abilities

68% (44 of 65) of patients on VYVGART achieved a significant improvement in their ability to perform daily activities*

Compared to 30% (19 of 64) of patients on placebo plus current treatment

Reduced muscle weakness

63% (41 of 65) of patients on VYVGART achieved a significant reduction in muscle weakness†

Compared to 14% (9 of 64) of patients on placebo plus current treatment

VYVGART was evaluated in a global study of adults with anti-AChR antibody positive gMG.

The study examined the safety and effectiveness of VYVGART in 167 adults (18 years or older) with gMG. In addition to their current treatment, patients received either VYVGART or placebo.

*Improvement maintained for 4 or more weeks was measured by a decrease of 2 or more points on the Myasthenia Gravis Activities of Daily Living (MG-ADL) scale, with the first reduction occurring no later than 1 week after the last infusion of treatment cycle 1. The MG-ADL scale assesses the impact of gMG on daily functions by measuring 8 signs or symptoms that are commonly affected in gMG. Each item is measured on a 4-point scale, where a score of 0 represents normal function and a score of 3 represents the loss of ability to perform that function. Total scores range from 0 to 24 points, with a higher score showing more severe gMG.

†Improvement maintained for 4 or more weeks was measured by a decrease of 3 or more points on the Quantitative Myasthenia Gravis (QMG) scale, with the first reduction occurring no later than 1 week after the last infusion of treatment cycle 1. The QMG scale assesses muscle weakness in gMG based on 13 items. Each item is assessed on a 4-point scale, where a score of 0 represents no muscle weakness and a score of 3 represents severe muscle weakness. Total scores range from 0 to 39, with a higher score meaning muscle weakness is more severe.

SELECT IMPORTANT SAFETY INFORMATION


VYVGART and VYVGART HYTRULO may cause serious side effects, including allergic reactions (hypersensitivity reactions). VYVGART and VYVGART HYTRULO can cause allergic reactions such as rashes, swelling under the skin, and shortness of breath. Hives were also observed in patients treated with VYVGART HYTRULO. Serious allergic reactions, such as trouble breathing and decrease in blood pressure leading to fainting have been reported with efgartigimod alfa-fcab.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

SAFETY

VYVGART was safe in treating most patients in the study and has a proven safety profile

In the study, the following side effects were reported in at least 5% of patients on VYVGART and more frequently than in patients on placebo



Most infections in patients on VYVGART for IV infusion were mild to moderate. Additionally, more patients on VYVGART for IV infusion vs. placebo had lower white blood cell counts that were mild to moderate in severity.

VYVGART for IV infusion may increase the risk of allergic reactions including rash, swelling under the skin, and shortness of breath. It may also increase the risk of serious allergic reactions such as trouble breathing and decrease in blood pressure leading to fainting, as well as infusion-related reactions including increase in blood pressure, chills, shivering, and chest, abdominal, and back pain. These reactions can occur during or after treatment and can cause doctors to pause or, in some cases, stop treatment.

AChR=acetylcholine receptor; gMG=generalized myasthenia gravis; IV=intravenous
Visit [VYVGART.com/glossary](https://www.vyvgart.com/glossary) for a glossary of terms.

➔ See more VYVGART study data at [VYVGART.com](https://www.vyvgart.com)

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).



ED & BREAKFAST

**GENERALIZED
MYASTHENIA
GRAVIS**
*isn't canceling this
weekend getaway*

VYVGART HYTRULO FOR SUBCUTANEOUS INJECTION



VYVGART Hytrulo is a ~30 to 90-second subcutaneous injection given at an infusion center, doctor's office, or at home*†

VYVGART Hytrulo contains 2 active ingredients: efgartigimod alfa, which is the same active ingredient in VYVGART for IV infusion, and hyaluronidase (human recombinant). Hyaluronidase helps to increase the distribution and absorption of ingredients into the body. For more information on what efgartigimod alfa does, see pages 6 and 7.

IV=intravenous

*For at least 30 minutes after your injection, a healthcare professional will monitor you for reactions.

†In some cases, VYVGART Hytrulo may also be given at home by a trained nurse.

SELECT IMPORTANT SAFETY INFORMATION

Do not use VYVGART if you have a serious allergy to efgartigimod alfa or any of the other ingredients in VYVGART. Do not use VYVGART HYTRULO if you have a serious allergy to efgartigimod alfa, hyaluronidase, or any of the other ingredients in VYVGART HYTRULO. VYVGART and VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

EFFECTIVENESS

VYVGART Hytrulo was found to be effective for adults with anti-AChR antibody positive gMG

The effectiveness of VYVGART Hytrulo was established through 2 studies.

In the VYVGART Hytrulo study, VYVGART Hytrulo for subcutaneous injection had a similar reduction in the harmful AChR antibodies that cause gMG symptoms as VYVGART for IV infusion.

In the VYVGART for IV infusion study, a reduction in harmful AChR antibodies was associated with improved daily abilities (reduction in total MG-ADL score).

See page 10 for study results.

VYVGART Hytrulo study design: The study compared antibody reduction in 110 adults (18 years or older) with gMG who remained on their current gMG treatment and were given either VYVGART Hytrulo for subcutaneous injection or VYVGART for IV infusion.



SELECT IMPORTANT SAFETY INFORMATION

VYVGART and VYVGART HYTRULO may cause serious side effects, including infusion-related reactions. VYVGART and VYVGART HYTRULO can cause infusion-related reactions. The most frequent symptoms and signs reported with efgartigimod alfa-fcab were high blood pressure, chills, shivering, and chest, abdominal, and back pain.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

SAFETY



VYVGART Hytrulo was safe in treating most patients studied

The safety of VYVGART Hytrulo was established through 2 studies.

The overall safety of VYVGART Hytrulo for subcutaneous injection was consistent with the safety profile of VYVGART for IV infusion, except patients on VYVGART Hytrulo for subcutaneous injection experienced more injection site reactions.

In the VYVGART for IV infusion study, the most common side effects for VYVGART for IV infusion-treated patients were respiratory tract infection, headache, and urinary tract infection. Additionally, more patients on VYVGART for IV infusion vs. placebo had lower white blood cell counts that were mild to moderate in severity.

See page 11 for study results.

In the VYVGART Hytrulo study, the most common side effects of VYVGART Hytrulo-treated patients were injection site reactions, 38% (21 of 55), and headache. Injection site reactions included injection site rash, redness of the skin, itching sensation, bruising, pain, and hives.

In the VYVGART Hytrulo study and its long-term safety study:

- All injection site reactions were mild to moderate and didn't lead to treatment discontinuation
- The majority of injection site reactions occurred within 24 hours of injection and resolved on their own
- The majority of injection site reactions occurred during the first treatment cycle and decreased with each subsequent cycle

VYVGART Hytrulo may increase the risk of allergic reactions including rash, swelling under the skin, shortness of breath, and hives. It may also increase the risk of serious allergic reactions such as trouble breathing and decrease in blood pressure leading to fainting, as well as infusion-related reactions including increase in blood pressure, chills, shivering, and chest, abdominal, and back pain. These reactions can occur during or after treatment and can cause doctors to pause or, in some cases, stop treatment.

AChR=acetylcholine receptor; gMG=generalized myasthenia gravis; IV=intravenous
Visit [VYVGART.com/glossary](https://www.vyvgart.com/glossary) for a glossary of terms.

➔ Visit [VYVGART.com](https://www.vyvgart.com) for more information


Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

DOSING SCHEDULE

VYVGART dosing—make the most of a treatment plan that's designed for you

VYVGART is given in treatment cycles with a personalized break between each cycle

Each treatment cycle is 4 treatments, 1 given each week.



You'll work with your doctor to personalize your breaks between cycles based on your symptoms and side effects.



When your symptoms start to return, talk to your doctor to help them decide when you need another cycle.

For example: Your doctor may start you with a 4-week break after each of your first 3 treatment cycles. Your doctor will continue to adjust your time between cycles, based on your symptoms and side effects.

The safety of starting the next treatment cycle sooner than 4 weeks from the last treatment of the previous treatment cycle has not been established. If you miss a scheduled treatment, VYVGART may be given up to 3 days after the scheduled treatment day.

SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART or VYVGART HYTRULO, tell your doctor if you: take any medicines, including prescription and non-prescription medicines, supplements, or herbal medicines; have received or are scheduled to receive a vaccine (immunization); or have any allergies or medical conditions, including if you are pregnant or planning to become pregnant, or are breastfeeding.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

Personalized breaks may help you get the most out of your treatment plan



PREZINA'S

break between cycles:

4 weeks



FRED'S

break between cycles:

6 weeks



DOUG'S

break between cycles:

9 weeks

Breaks between cycles vary based on each patient's symptoms and side effects. Learn about real patients' breaks between cycles on VYVGARTDosing.com.

Tracking your symptoms consistently is important

Starting your next treatment cycle is based on your doctor's evaluation of your symptoms and any side effects you may experience. Use the MG-ADL scale to help track your symptoms.

Prezina, Fred, and Doug are paid spokespeople of argenx.
MG-ADL=Myasthenia Gravis Activities of Daily Living

➔ Download the doctor discussion guide at VYVGARTDiscussionGuide.com to get your MG-ADL score and more conversation starters

SELECT IMPORTANT SAFETY INFORMATION

Do not use VYVGART if you have a serious allergy to efgartigimod alfa or any of the other ingredients in VYVGART. Do not use VYVGART HYTRULO if you have a serious allergy to efgartigimod alfa, hyaluronidase, or any of the other ingredients in VYVGART HYTRULO. VYVGART and VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

**SUPPORT FROM
MY VYVGART PATH**

My VYVGART[®] Path

Here for you during your VYVGART treatment journey

My VYVGART Path is a **Patient Support Program** that provides personalized support from a Nurse Case Manager and committed support team. They are dedicated to listening to you and answering questions you may have about VYVGART during your treatment journey.



➔ To get started, ask your neurologist to enroll you in My VYVGART Path or visit myVYVGARTpath.com to learn more




Understand your VYVGART treatment

A Nurse Case Manager can provide educational information to help you learn about VYVGART for IV infusion and VYVGART Hytrulo for subcutaneous injection and understand the infusion and injection processes.



Feel empowered with resources and information

Nurse Case Managers can support you throughout your VYVGART treatment journey to help you in following your doctor's treatment plan.



Navigate the insurance process

You may have questions about your insurance coverage, want to know more about the cost of treatment, or just need information about your out-of-pocket costs for VYVGART. With My VYVGART Path, you'll find support in understanding each step of the insurance process.



Understand potential financial assistance programs

If you have financial concerns or gaps in insurance coverage for your VYVGART therapy, a Nurse Case Manager is here for you with information and support.

Please see **Important Safety Information** throughout. Please see full **Prescribing Information** for **VYVGART for IV infusion** and **VYVGART HYTRULO for subcutaneous injection**.

FREQUENTLY ASKED QUESTIONS

How is VYVGART given?

VYVGART is available as VYVGART for IV infusion or as VYVGART Hytrulo for subcutaneous injection. An IV infusion is given by inserting a needle into a vein so that medicine immediately enters the bloodstream. A subcutaneous injection delivers medicine into the layer beneath the skin.

Is VYVGART intended to be taken with other medications?

Yes, in the studies, patients stayed on their current gMG treatment while receiving VYVGART for IV infusion or VYVGART Hytrulo for subcutaneous injection. Talk to your doctor about the medications you are taking as part of your treatment plan.

What are the most common side effects of VYVGART?

Out of 84 patients in the study treated with VYVGART for IV infusion, the most common side effects included respiratory tract infection (33%), headache (32%), urinary tract infection (10%), tingling sensation (7%), and muscle pain (6%).* Most infections in patients on VYVGART for IV infusion

were mild to moderate. Additionally, more patients on VYVGART for IV infusion vs. placebo had lower white blood cell counts that were mild to moderate in severity.

The overall safety of VYVGART Hytrulo for subcutaneous injection was consistent with the safety profile of VYVGART for IV infusion, except patients on VYVGART Hytrulo for subcutaneous injection experienced more injection site reactions.

In the VYVGART Hytrulo study, the most common side effects of VYVGART Hytrulo-treated patients were injection site reactions, 38% (21 of 55), and headache. Injection site reactions included injection site rash, redness of the skin, itching sensation, bruising, pain, and hives.

In the VYVGART Hytrulo study and its long-term safety study, all injection site reactions were mild to moderate and didn't lead to treatment discontinuation. The majority occurred within 24 hours of injection and resolved on their own. The majority occurred during the first treatment cycle and decreased with each subsequent cycle.

*In ≥5% treated with VYVGART and more frequently than placebo.

SELECT IMPORTANT SAFETY INFORMATION

Tell your doctor if you have signs or symptoms of an infection, allergic reaction, or infusion-related reaction. These can happen while you are receiving your VYVGART or VYVGART HYTRULO treatment or afterward. Your doctor may need to pause or stop your treatment. Contact your doctor immediately if you have signs or symptoms of a serious allergic reaction.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

Do I have to get any vaccines before taking VYVGART?

Some types of vaccines are not recommended while taking VYVGART. Talk to your doctor about vaccines, including when you need to receive age-appropriate vaccines, before starting a treatment cycle of VYVGART.

How many patients in the VYVGART for IV infusion study continued to the extended safety study?

91% of the patients in the study who were eligible chose to enter a 3-year extension safety study where they would receive VYVGART for IV infusion.

What should I do if I miss a treatment?

If you miss an IV infusion or subcutaneous injection, VYVGART may be given up to 3 days after the scheduled treatment day. Work with your neurologist to reschedule your next treatment.

How do I track my symptoms?

Download the MG-ADL assessment tool on VYVGART.com to start tracking your symptoms. If you are enrolled in My VYVGART Path, a physical journal will be mailed to you and a Nurse Case Manager can show you how to use it as you get started.

How do I join the My VYVGART Path program?

After you and your neurologist decide on VYVGART, your doctor's office will be able to enroll you in My VYVGART Path. Then, a Nurse Case Manager will call, welcome you to the program, and offer assistance as you get started on your treatment journey with VYVGART for IV infusion or VYVGART Hytrulo for subcutaneous injection.

IV=intravenous

Visit [VYVGART.com/glossary](https://www.vyvgart.com/glossary) for a glossary of terms.

 **More information available at [VYVGART.com](https://www.vyvgart.com)**

SELECT IMPORTANT SAFETY INFORMATION

Before taking VYVGART or VYVGART HYTRULO, tell your doctor if you: take any medicines, including prescription and non-prescription medicines, supplements, or herbal medicines; have received or are scheduled to receive a vaccine (immunization); or have any allergies or medical conditions, including if you are pregnant or planning to become pregnant, or are breastfeeding.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

IMPORTANT SAFETY INFORMATION

Do not use VYVGART if you have a serious allergy to efgartigimod alfa or any of the other ingredients in VYVGART. Do not use VYVGART HYTRULO if you have a serious allergy to efgartigimod alfa, hyaluronidase, or any of the other ingredients in VYVGART HYTRULO. VYVGART and VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

VYVGART and VYVGART HYTRULO may cause serious side effects, including:

- **Infection.** VYVGART and VYVGART HYTRULO may increase the risk of infection. The most common infections for efgartigimod alfa-fcab-treated patients were urinary tract and respiratory tract infections. Signs or symptoms of an infection may include fever, chills, frequent and/or painful urination, cough, pain and blockage of nasal passages/sinus, wheezing, shortness of breath, fatigue, sore throat, excess phlegm, nasal discharge, back pain, and/or chest pain.
- **Allergic Reactions (hypersensitivity reactions).** VYVGART and VYVGART HYTRULO can cause allergic reactions such as rashes, swelling under the skin, and shortness of breath. Hives were also observed in patients treated with VYVGART HYTRULO. Serious allergic reactions, such as trouble breathing and decrease in blood pressure leading to fainting have been reported with efgartigimod alfa-fcab.
- **Infusion-Related Reactions.** VYVGART and VYVGART HYTRULO can cause infusion-related reactions. The most frequent symptoms and signs reported with efgartigimod alfa-fcab were high blood pressure, chills, shivering, and chest, abdominal, and back pain.

Tell your doctor if you have signs or symptoms of an infection, allergic reaction, or infusion-related reaction. These can happen while you are receiving your VYVGART or VYVGART HYTRULO treatment or afterward. Your doctor may need to pause or stop your treatment. Contact your doctor immediately if you have signs or symptoms of a serious allergic reaction.

Before taking VYVGART or VYVGART HYTRULO, tell your doctor if you:

- take any medicines, including prescription and non-prescription medicines, supplements, or herbal medicines,
- have received or are scheduled to receive a vaccine (immunization), or
- have any allergies or medical conditions, including if you are pregnant or planning to become pregnant, or are breastfeeding.

What are the common side effects of VYVGART and VYVGART HYTRULO?

The most common side effects in efgartigimod-alfa-fcab-treated patients were respiratory tract infection, headache, and urinary tract infection. Additional common side effects with VYVGART HYTRULO are injection site reactions, including rash, redness of the skin, itching sensation, bruising, pain, and hives.

These are not all the possible side effects of VYVGART and VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to the US Food and Drug Administration at 1-800-FDA-1088.

What is VYVGART® (efgartigimod alfa-fcab) for intravenous (IV) infusion and what is VYVGART® HYTRULO (efgartigimod alfa and hyaluronidase-qvfc) for subcutaneous injection?

VYVGART and VYVGART HYTRULO are both prescription medicines, each used to treat a condition called generalized myasthenia gravis, which causes muscles to tire and weaken easily throughout the body, in adults who are positive for antibodies directed toward a protein called acetylcholine receptor (anti-AChR antibody positive).

Please see the full Prescribing Information for VYVGART and the full Prescribing Information for VYVGART HYTRULO, and talk to your doctor.



VYVGART® Hytrulo
(efgartigimod alfa and
hyaluronidase-qvfc)
Subcutaneous Injection
180 mg/mL and 2000 U/mL vial

VYVGART®
(efgartigimod alfa-fcab)
Injection for Intravenous Use
400 mg/20 mL vial

Picture your life in motion

- ✓ When added to their current gMG treatment, VYVGART for IV infusion helped most patients in the study with anti-AChR antibody positive gMG achieve **improved daily abilities***
- ✓ VYVGART Hytrulo **was found to be effective** for adults with anti-AChR antibody positive gMG
- ✓ VYVGART for IV infusion and VYVGART Hytrulo for subcutaneous injection were **safe in treating most patients studied**

Talk to your neurologist to see if VYVGART is right for you



Use the camera on your smartphone to scan the QR code and download your VYVGART Doctor Discussion Guide.

*Improvement maintained for 4 or more weeks was measured by a decrease of 2 or more points on the Myasthenia Gravis Activities of Daily Living (MG-ADL) scale, with the first reduction occurring no later than 1 week after the last infusion of treatment cycle 1. The MG-ADL scale assesses the impact of gMG on daily functions by measuring 8 signs or symptoms that are commonly affected in gMG. Each item is measured on a 4-point scale, where a score of 0 represents normal function and a score of 3 represents the loss of ability to perform that function. Total scores range from 0 to 24 points, with a higher score showing more severe gMG.

AChR=acetylcholine receptor; gMG=generalized myasthenia gravis; IV=intravenous
Visit [VYVGART.com/glossary](https://www.vyvgart.com/glossary) for a glossary of terms.

➔ Call 1-833-VYVGART (1-833-898-4278) or visit [VYVGART.com](https://www.vyvgart.com) to find out more

SELECT IMPORTANT SAFETY INFORMATION

VYVGART and VYVGART HYTRULO may increase the risk of infection. The most common infections for efgartigimod alfa-fcab-treated patients were urinary tract and respiratory tract infections.

Please see Important Safety Information throughout. Please see full Prescribing Information for [VYVGART for IV infusion](#) and [VYVGART HYTRULO for subcutaneous injection](#).

VYVGART is a registered trademark of argenx. VYVGART Hytrulo is a trademark of argenx.
For U.S. audiences only. ©2024 argenx US-ESC-22-00011 V2 05/2024

argenx 